

Ch. 1.6 Methods of Proofs

The following methods of proofs are discussed in our text:

- Vacuous proof
 - Trivial proof
 - Direct proof
 - Indirect proof
 - Proof by contradiction
 - Proof by cases.
- A **vacuous proof** of an implication happens when the hypothesis of the implication is always false.

Example 1: Prove that if x is a positive integer and $x = -x$, then $x^2 = x$.
proof.

- An implication is **trivially true** when its conclusion is always true.
- A declared mathematical proposition whose truth-value is unknown is called a **conjecture**.

One of the main functions of a mathematician (and a computer scientist) is to decide the truth-value of their claims (or someone else's claims).

- If a conjecture is proven true we call it a *theorem*, *lemma* or *corollary*; if it proven false, then usually discarded.
- A **proof** is a sequence of statements bound together by the rules of logic, definitions, previously proven theorems, simple algebra and axioms.

Definition: An integer n is even if there exists an integer k such that $n = 2k$. An integer n is odd if there exists an integer k such that $n = 2k + 1$.

Axiom (Closure of addition over the integers): If a and b are integers, then $a + b$ is an integer.

Axiom (Closure of multiplication over the integers): If a and b are integers, then $a \cdot b$ is an integer.

Example 2: Prove or disprove that the product of two irrational numbers is irrational.

Example 3: prove lemma 1.

Lemma 1. If n is even, then n^2 is even.

Proof:

Example 4: Prove lemma 2

Lemma 2. If n^2 is even, then n is even.

Proof:

Theorem 1: An integer n is even if and only if n^2 is even.

Proof: If n is even, then n^2 is even is true by Lemma 1. The converse, if n^2 is even, then n is even is true by Lemma 2. Hence the biconditional statement n is even if and only if n^2 is even is true. $\square$

Example 5: Prove that the sum of two odd integers is even. i.e. If p and q are odd integers, then $p + q$ is an even integer.

Proof:

Summary. If we are proving $p \rightarrow q$, then

A direct proof begins by assuming p is true.	An indirect proof begins by assuming $\sim q$ is true.
$\vdots$	$\vdots$
$\vdots$	$\vdots$
until we conclude q .	until we conclude $\sim p$.

Example 6: Prove that $\sqrt{2}$ is irrational.

Proof: Assume by way of contradiction that can be represented as a quotient of two integers p/q where q is not zero. Further, we assume that p/q is in lowest terms, i.e. we assume that

The integers p and q have no common factor. (1)

Thus, by assumption $\sqrt{2} = p/q$, and now squaring both sides yields

$$2 = \frac{p^2}{q^2} \quad \text{or} \quad p^2 = 2q^2 \quad (2)$$

This implies that p^2 is even, and by Theorem 1, p must also be even. So we write $p = 2k$, substitute into the second equation of (2), and by cancellation we see that

$$q^2 = 2k^2. \quad (3)$$

This says that q^2 is even, and again by Theorem 1, q must also be even. From statements (2) and (3), it follows that

$$p \text{ and } q \text{ both have } 2 \text{ as a common factor.} \quad (4)$$

Statements (1) and (4) are contradictory. Thus, $\sqrt{2}$ is not rational.



Summary. If we are proving $p \rightarrow q$, then

A direct proof begins by assuming p is true. : : until we conclude q .	An indirect proof begins by assuming $\sim q$ is true. : : until we conclude $\sim p$.	An proof by contradiction begins by assuming $p \wedge \sim q$ is true. : : until we reach a contradiction
---	--	---

Example 7: Prove that if $3n + 2$ is odd, then n is odd.

- i. Write the proposition in symbolic logic notation.
- ii. Write the negation of the proposition in symbolic logic notation.
- iii. *Proof:*

Example 8: Prove that there is an even prime number.

Definition. Let x be a real number. Then $|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$.

Example 9: Prove if x is a real number, then $|-x| = |x|$.

Definition. A function $f:A \rightarrow B$ is *one-to-one* if and only if

$$\forall x \forall y (f(x) = f(y) \rightarrow x = y),$$

which is logically equivalent to its contrapositive

$$\forall x \forall y (x \neq y \rightarrow f(x) \neq f(y)).$$

Example 10: Prove that the real valued function $f(x) = x + I$ is one-to-one.